

# Non-surgical treatment of pulpitis, including those with history of spontaneous pain, using a combination of antibacterial drugs

Toyohiko Takushige<sup>1,3</sup>, D.D.S., Ph.D., Edward Venzon Cruz<sup>1,2</sup>, D.D.S., Ph.D., Md. Ali Asgor Moral<sup>1,2</sup>, D.D.S., Ph.D., Etsuro Hoshino<sup>1,2</sup>, D.D.S., Ph.D.

Oral Ecology in Health and Infection<sup>1</sup> and Cariology Research Unit<sup>2</sup>,  
Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan,  
and Takushige Dental Clinic<sup>3</sup>, Sendai, Japan

---

## Abstract

This is a retrospective clinical study of 360 teeth diagnosed with pulpitis and treated locally with a combination of three antibacterial drugs (ciprofloxacin, metronidazole and minocycline, “3Mix-MP”) without a pulpectomy procedure. Consecutive patients with a clinical diagnosis of pulpitis had pre-operative factors collected (spontaneous pain, pulp exposure, depth of carious lesion) and were treated by placement of 3Mix-MP onto the pulpal floor of the carious lesion where softened dentin was intentionally left, if any was present. The treated lesions were then sealed by glass-ionomer cement and restored by resin inlays. A good clinical outcome was defined as the lack of any spontaneous pain, no mechanical allodynia to biting and the presence of pulpal responsiveness to cold or electrical stimuli. Using these criteria, a good clinical outcome was found in 342 (95%) of the 360 cases. Six cases progressed to pulpal necrosis and the remaining 12 cases required re-treatment using the same 3Mix-MP, resulting in a subsequent good outcome. Recalcification of softened dentin was evident on postoperative radiographs. These data suggest that 3Mix-MP may be worth evaluating in prospective randomized clinical trials for treatment of pulpitis including cases of so-called “irreversible” pulpitis.

---

## Key words:

3Mix-MP, LSTR 3Mix-MP, SavePulp Therapy, pulpitis with spontaneous pain

## Correspondent:

Professor HOSHINO Etsuro

Oral Ecology in Health and Infection,

Niigata University Graduate School of Medical and Dental Sciences

Gakkocho-dori 2, Niigata 951-8514 Japan

Fax: +81 (0)25-227-0806; Tel: +81 (0)25-227-2838; e-mail: hoshino@dent.niigata-u.ac.jp

---

Supported by the Grants-in-aid for Scientific Research (14406028, and 17390500) from the Ministry of Education, Culture, Sports, Science and Technology, and by the grants for the Joint Research Program from the Japan Society for the Promotion of Science.

---